

# 3M™ Tegaderm™ Foam High Performance Foam Non-Adhesive Dressing

## Product Description

3M™ Tegaderm™ High Performance Foam Non-Adhesive Dressing is a highly absorbent, breathable, non-adherent wound dressing. It is constructed from a conformable polyurethane foam pad, absorbent nonwoven layers, and a top layer of printed adhesive film. This waterproof film is moisture vapor permeable which prevents wound exudate strike-through and acts as a barrier to outside contamination, including bacteria and viruses.\* The dressing maintains a moist wound environment, which has been shown to enhance wound healing.

## Indications for Use

Tegaderm™ Foam Dressing is indicated for use as a primary or secondary dressing for low- to highly-exuding partial and full thickness wounds such as pressure ulcers, venous leg ulcers, neuropathic ulcers, arterial ulcers, skin tears, surgical wounds, abrasions, superficial partial thickness burns and donor sites. The dressing is suitable for use with compression therapy.

The dressing is not designed, sold or intended for use except as indicated. Tegaderm™ Foam Dressing is not intended for use in pressure reduction or as a surgical sponge.

## Contraindications

None Known

## Warnings

None Known

## Precautions


1. Do not use Tegaderm™ Foam Dressing in combination with oxidizing agents such as hydrogen peroxide or hypochlorite solution. These solutions may affect the structure and performance of the polyurethane foam.
2. At the time of dressing change, if the dressing is adhered to the wound surface, saturate with normal saline and gently loosen.
3. Tegaderm™ Foam Dressing may be used on infected wounds under the care of a health care professional. If the wound does not begin to show signs of healing or if any other unexpected symptoms occur, consult a health care professional.

## Directions for Use

Note: Follow facility guidelines for infection control.

1. Cleanse wound and surrounding skin according to facility policy.
2. Remove dressing from package. (Figure 1)
3. Determine appropriate size dressing needed to ensure that the foam pad overlaps the wound edges. The dressing may be cut as necessary to accommodate wound size and contours. Use of non-sterile product may increase the risk of contamination and infection. (Figure 2)
4. Position the dressing, printed film side up, over the wound with edges overlapping onto intact skin. Secure with elastic or cohesive wrap, tape, or other methods. (Figure 3)

Note: Covering the entire foam dressing with an occlusive tape may reduce the breathability of the film backing. To maintain moisture vapor exchange, tape only edges of the dressing in picture frame manner.



## Dressing Change Considerations

Tegaderm™ Foam Dressing may remain in place for up to seven days. As the dressing absorbs, exudate will wick to the top of the dressing, and discoloration may be noticeable. When the exudate spreads to the edges of the dressing or the dressing begins to leak, a dressing change is indicated. Dressing change frequency will depend on the type of wound, volume of exudate, and clinical situation. Change at least every seven days, or as indicated per treatment protocol.

## Storage

Store at room temperature.

## How Supplied

Supplied in individually packaged sterile dressings. Product is sterile until package is damaged or opened.

\* In vitro testing shows that the transparent film provides a viral barrier from viruses 27 nm in diameter or larger while the dressing remains intact without leakage.

## Explanation of Symbols

 Do not use if package is damaged or open

 This product and package do not contain natural rubber latex.

 Caution, see instructions for use

 Do not reuse

 Use by date

 Batch code

 Manufacturer

 Sterilized using irradiation

**CE 0086**

Made in U.S.A. by

**3M Health Care**

2510 Conway Ave.

St. Paul, MN 55144

1-800-228-3957

[www.3M.com/Tegaderm](http://www.3M.com/Tegaderm)

3M and Tegaderm are trademarks of 3M.

Used under license in Canada.

© 2013, 3M. All rights reserved.

Issue Date: 2013-04

34-8712-6614-3

**EC REP 3M Deutschland GmbH**

**Health Care Business**

Carl-Schurz-Str. 1

41453 Neuss

Germany

3M Canada Company

P.O. Box / C.P. 5757 London, Ontario, N6A 4T1

1-800-364-3577

3M et Tegaderm sont des marques de commerce de 3M, utilisées sous licence au Canada.

© 2013, 3M. Tous droits réservés.