

3M ESPE

Single Bond Universal

Single Bond Universal DCA

- en** Adhesive / Dual Cure Activator
- ru** Адгезив / Активатор для двойного твердения
- csk** Adhesive / Dvojnomomenizační aktivátor
- hr** Adheziv / Aktivator za dvostruko stvrdnjavanje
- hu** Ragasztóanyag / Kettős kőtésű aktivátor
- pl** Klej / Podwójnie utwardzający aktywator
- de** Adeziv / Aktivator de polymerizare dubă
- sk** Lepidlo / Aktivátor podvojného tvrdnutia

Instructions for Use
Инструкции по применению
Указания за употреба
Ирире за употребу

3M Deutschland GmbH
Dental Products
Carl-Schurz-Str. 1
4493 Neuss - Germany

3M, ESPE, Clinpro и RelyX are trademarks of 3M or
3M Deutschland GmbH. © 2013 3M. All rights reserved.

Felhasználási tudnivaló
Instrukcja użycia
Instructiuni de utilizare
Informácie pre požívanie



44000763977/03

Instructions for Use
Инструкции по применению
Указания за употреба
Ирире за употребу

3M Deutschland GmbH
Dental Products
Carl-Schurz-Str. 1
4493 Neuss - Germany

3M, ESPE, Clinpro и RelyX are trademarks of 3M or
3M Deutschland GmbH. © 2013 3M. All rights reserved.

Felhasználási tudnivaló
Instrukcja użycia
Instructiuni de utilizare
Informácie pre požívanie



44000763977/03

Instructions for Use
Инструкции по применению
Указания за употреба
Ирире за употребу

3M Deutschland GmbH
Dental Products
Carl-Schurz-Str. 1
4493 Neuss - Germany

3M, ESPE, Clinpro и RelyX are trademarks of 3M or
3M Deutschland GmbH. © 2013 3M. All rights reserved.

Felhasználási tudnivaló
Instrukcja użycia
Instructiuni de utilizare
Informácie pre požívanie



44000763977/03

Instructions for Use
Инструкции по применению
Указания за употреба
Ирире за употребу

3M Deutschland GmbH
Dental Products
Carl-Schurz-Str. 1
4493 Neuss - Germany

3M, ESPE, Clinpro и RelyX are trademarks of 3M or
3M Deutschland GmbH. © 2013 3M. All rights reserved.

Felhasználási tudnivaló
Instrukcja użycia
Instructiuni de utilizare
Informácie pre požívanie



44000763977/03

Instructions for Use
Инструкции по применению
Указания за употреба
Ирире за употребу

3M Deutschland GmbH
Dental Products
Carl-Schurz-Str. 1
4493 Neuss - Germany

3M, ESPE, Clinpro и RelyX are trademarks of 3M or
3M Deutschland GmbH. © 2013 3M. All rights reserved.

Felhasználási tudnivaló
Instrukcja użycia
Instructiuni de utilizare
Informácie pre požívanie



44000763977/03

Instructions for Use
Инструкции по применению
Указания за употреба
Ирире за употребу

3M Deutschland GmbH
Dental Products
Carl-Schurz-Str. 1
4493 Neuss - Germany

3M, ESPE, Clinpro и RelyX are trademarks of 3M or
3M Deutschland GmbH. © 2013 3M. All rights reserved.

Felhasználási tudnivaló
Instrukcja użycia
Instructiuni de utilizare
Informácie pre požívanie



44000763977/03

Rinse thoroughly with water and dry with water-free and oil-free air or with cotton pellets; do not overdry.

Root Surface Desensitization

Clean the root surface as usual (e.g., with polishing paste or powder).

Then rinse off the veneer with RelyX® Veneer Cement following the manufacturer's instructions for restorative procedures.

After rinsing, apply Single Bond Universal Adhesive.

Pre-treatment of Fillings or Restorations During Repair

In cases of restorative procedures on fillings or restorations, remove any excess from the veneer.

We recommend light-curing the gingival edges first, then the middle and the incisal edges. Avoid eye contact with the light source.

Procedure for a composite filling or restoration

In cases of restorative procedures on fillings or restorations, remove any excess from the veneer.

We recommend light-curing the gingival edges first, then the middle and the incisal edges. Avoid eye contact with the light source.

After rinsing, apply Single Bond Universal Adhesive.

Dosing from the Bottle

The dosing bottle contains a thumb depression so that it can be opened to one side. The cap will remain open if the opening angle is > 90°.

To open, take the bottle in your hand, place your thumb in the thumb depression and push the cap to the side until it remains in the open position.

If necessary, repeat the dispensing procedure for the remaining teeth.

Subsequently direct a gentle stream of air over the liquid for about 5 sec until no longer needed and the solvent has evaporated completely.

Use a available light-curing device to polymerize the adhesive for 10 sec on every surface of the filling.

When the film has been completely cured, grind down the occlusion and sand the margins of the filling to the ground plane to protect the filling from drying out.

After curing, use a damp cloth or cotton pellet to remove the oxygen inhibition layer.

Hygiene and Disinfection

Clean any contaminated adhesive and activator bottles with commonly used agents and disinfect them in accordance with the commonly used hygiene regulations.

Note:

Single Bond Universal can prematurely polymerize when exposed to natural or artificial lighting. Avoid intensive light exposure during application.

Storage and Stability

Store Single Bond Universal and Single Bond Universal DCA at 0-25°C/ 36-77°F.

Do not use after the expiration date.

Customer Information

No person is authorized to provide any information which deviates from the information provided in this instruction sheet.

Dosing Single Bond Universal DCA Activator

For each application of the adhesive, the dispensing applicator can be bent with the thumb in the L-Pop® finger slot, then only pull the dispensing applicator out of the finger slot far enough so that the narrow section of the dispensing applicator shaft can be seen. Fold the applicator shaft at this location over the thumb.

Hygiene and Disinfection

Clean any contaminated adhesive and activator bottles with commonly used agents and disinfect them in accordance with the commonly used hygiene regulations.

Note:

Single Bond Universal DCA activates the dual curing mechanism of Single Bond Universal and is mixed with the adhesive under the same conditions.

In a "self-curing" procedure to enable the shortest possible treatment time, the adhesive must be applied directly to the tooth enamel and to minimize post-operative sensitivity.

In a "total etching" procedure with a prior phosphoric acid etching step, e.g., if the etching cannot be restricted to the tooth enamel.

Single Bond Universal DCA activates the dual curing mechanism of Single Bond Universal and is mixed with the adhesive under the same conditions.

In a "self-curing" procedure to enable the shortest possible treatment time, the adhesive must be applied directly to the tooth enamel and to minimize post-operative sensitivity.

In a "total etching" procedure with a prior phosphoric acid etching step, e.g., if the etching cannot be restricted to the tooth enamel.

Single Bond Universal DCA activates the dual curing mechanism of Single Bond Universal and is mixed with the adhesive under the same conditions.

In a "self-curing" procedure to enable the shortest possible treatment time, the adhesive must be applied directly to the tooth enamel and to minimize post-operative sensitivity.

In a "total etching" procedure with a prior phosphoric acid etching step, e.g., if the etching cannot be restricted to the tooth enamel.

Single Bond Universal DCA activates the dual curing mechanism of Single Bond Universal and is mixed with the adhesive under the same conditions.

In a "self-curing" procedure to enable the shortest possible treatment time, the adhesive must be applied directly to the tooth enamel and to minimize post-operative sensitivity.

In a "total etching" procedure with a prior phosphoric acid etching step, e.g., if the etching cannot be restricted to the tooth enamel.

Single Bond Universal DCA activates the dual curing mechanism of Single Bond Universal and is mixed with the adhesive under the same conditions.

In a "self-curing" procedure to enable the shortest possible treatment time, the adhesive must be applied directly to the tooth enamel and to minimize post-operative sensitivity.

In a "total etching" procedure with a prior phosphoric acid etching step, e.g., if the etching cannot be restricted to the tooth enamel.

Single Bond Universal DCA activates the dual curing mechanism of Single Bond Universal and is mixed with the adhesive under the same conditions.

In a "self-curing" procedure to enable the shortest possible treatment time, the adhesive must be applied directly to the tooth enamel and to minimize post-operative sensitivity.

In a "total etching" procedure with a prior phosphoric acid etching step, e.g., if the etching cannot be restricted to the tooth enamel.

Single Bond Universal DCA activates the dual curing mechanism of Single Bond Universal and is mixed with the adhesive under the same conditions.

In a "self-curing" procedure to enable the shortest possible treatment time, the adhesive must be applied directly to the tooth enamel and to minimize post-operative sensitivity.

In a "total etching" procedure with a prior phosphoric acid etching step, e.g., if the etching cannot be restricted to the tooth enamel.

Single Bond Universal DCA activates the dual curing mechanism of Single Bond Universal and is mixed with the adhesive under the same conditions.

In a "self-curing" procedure to enable the shortest possible treatment time, the adhesive must be applied directly to the tooth enamel and to minimize post-operative sensitivity.

In a "total etching" procedure with a prior phosphoric acid etching step, e.g., if the etching cannot be restricted to the tooth enamel.

Single Bond Universal DCA activates the dual curing mechanism of Single Bond Universal and is mixed with the adhesive under the same conditions.

In a "self-curing" procedure to enable the shortest possible treatment time, the adhesive must be applied directly to the tooth enamel and to minimize post-operative sensitivity.

In a "total etching" procedure with a prior phosphoric acid etching step, e.g., if the etching cannot be restricted to the tooth enamel.

Single Bond Universal DCA activates the dual curing mechanism of Single Bond Universal and is mixed with the adhesive under the same conditions.

In a "self-curing" procedure to enable the shortest possible treatment time, the adhesive must be applied directly to the tooth enamel and to minimize post-operative sensitivity.

In a "total etching" procedure with a prior phosphoric acid etching step, e.g., if the etching cannot be restricted to the tooth enamel.

Single Bond Universal DCA activates the dual curing mechanism of Single Bond Universal and is mixed with the adhesive under the same conditions.

In a "self-curing" procedure to enable the shortest possible treatment time, the adhesive must be applied directly to the tooth enamel and to minimize post-operative sensitivity.

In a "total etching" procedure with a prior phosphoric acid etching step, e.g., if the etching cannot be restricted to the tooth enamel.

Single Bond Universal DCA activates the dual curing mechanism of Single Bond Universal and is mixed with the adhesive under the same conditions.

In a "self-curing" procedure to enable the shortest possible treatment time, the adhesive must be applied directly to the tooth enamel and to minimize post-operative sensitivity.

In a "total etching" procedure with a prior phosphoric acid etching step, e.g., if the etching cannot be restricted to the tooth enamel.

Single Bond Universal DCA activates the dual curing mechanism of Single Bond Universal and is mixed with the adhesive under the same conditions.

In a "self-curing" procedure to enable the shortest possible treatment time, the adhesive must be applied directly to the tooth enamel and to minimize post-operative sensitivity.

In a "total etching" procedure with a prior phosphoric acid etching step, e.g., if the etching cannot be restricted to the tooth enamel.

Single Bond Universal DCA activates the dual curing mechanism of Single Bond Universal and is mixed with the adhesive under the same conditions.

In a "self-curing" procedure to enable the shortest possible treatment time, the adhesive must be applied directly to the tooth enamel and to minimize post-operative sensitivity.

In a "total etching" procedure with a prior phosphoric acid etching step, e.g., if the etching cannot be restricted to the tooth enamel.

Single Bond Universal DCA activates the dual curing mechanism of Single Bond Universal and is mixed with the adhesive under the same conditions.

In a "self-curing" procedure to enable the shortest possible treatment time, the adhesive must be applied directly to the tooth enamel and to minimize post-operative sensitivity.

In a "total etching" procedure with a prior phosphoric acid etching step, e.g., if the etching cannot be restricted to the tooth enamel.

Single Bond Universal DCA activates the dual curing mechanism of Single Bond Universal and is mixed with the adhesive under the same conditions.

In a "self-curing" procedure to enable the shortest possible treatment time, the adhesive must be applied directly to the tooth enamel and to minimize post-operative sensitivity.

In a "total etching" procedure with a prior phosphoric acid etching step, e.g., if the etching cannot be restricted to the tooth enamel.

Single Bond Universal DCA activates the dual curing mechanism of Single Bond Universal and is mixed with the adhesive under the same conditions.

In a "self-curing" procedure to enable the shortest possible treatment time, the adhesive must be applied directly to the tooth enamel and to minimize post-operative sensitivity.

In a "total etching" procedure with a prior phosphoric acid etching step, e.g., if the etching cannot be restricted to the tooth enamel.

Single Bond Universal DCA activates the dual curing mechanism of Single Bond Universal and is mixed with the adhesive under the same conditions.

In a "self-curing" procedure to enable the shortest possible treatment time, the adhesive must be applied directly to the tooth enamel and to minimize post-operative sensitivity.

In a "total etching" procedure with a prior phosphoric acid etching step, e.g., if the etching cannot be restricted to the tooth enamel.

3M ESPE**Single Bond Universal****Single Bond Universal DCA**

s) Adhezív / Aktivátor dvojného střívání

c) Adhezívum / Aktivátoru duálního vytvárování

tr) Adhesive / Dual Cure Activator

et) Adhesiv / Kaksikäytöistä aktivator

lv) Adhezív / Divkárásának aktivátor

it) Adesiva / Dvugivoce klenitimo aktivator

uk) Adhesiv / Aktivátor podávajúci zaveterejníka

el) Συγκόλλητικό παράγοντα / Επερυποτητής για διπλό πολυμερέσιο

Navodila za uporabu
Informacije po upravljanju
Kullanma Kilavuzu
Kasutusuhend

3M Deutschland GmbH

Dental Products

44566 St. Ingbert - Germany

3M, ESPE, Clinpro and RelV-X are trademarks of 3M or 3M Deutschland GmbH. © 2013, 3M. All rights reserved.



0123



44000763977/03

Lietotāšanas informācija

Vartojimo instrukcija

Istoričkis & Viskoprostřítná

Plnoprropočes rýptón

CE

0123

44000763977/03

Temeljito izpirazite z vodo in posušite z zrakom brez primesi vode ali olja ali z vanimi kroglicami; ne susište preveč.

Postopek totalnega jedkanja

Uporabite običajen gel za jedkanje s fosforno kislino (pribilno 35 %), npr. Scotchbond Universal Echant, ki ga nanesete na sklenko in dentin, in posušite z vodo, potem da reakcija konča.

Uporabite običajen gel za jedkanje s fosforno kislino (pribilno 35 %), npr. Scotchbond Universal Echant, ki ga nanesete na pripravljeno in nepravljeno keramiko v sklopu sklenke in dentina, in posušite z vodo, potem da reakcija konča.

Temeljito izpirazite z vodo in posušite z zrakom brez primesi vode ali olja ali z vanimi kroglicami; ne susište preveč.

Desenzibilizacija površine keramike

Površina keramike je delitvena, kar občajno (npr. s polirno pasti ali peskanjem), temeljito izpirazite z vodo in posušite z zrakom brez primesi vode ali olja ali z vanimi kroglicami; ne susište preveč.

Nadključite z namenjenim adhezivom Single Bond Universal.

Predhodna poliranja zatviri restavracij med uporabami

Steklene keramike z keramiko, cirkonijem, keramike iz aluminijskega oksiida, kovino (krm. z amalgamom).

Temeljito delovite površine restavracije, ki jo želite lepliti, najbolje s posebnimi sponzni sredstvi za zdravje zraka in vodo.

Temeljito očistite s pršenjem vode in posušite z zrakom brez primesi vode ali olja ali z vanimi kroglicami.

Takoj nadaljujte z namenjenim adhezivom Single Bond Universal. Uporaba dodatne prima je potrebna.

Uporabite običajen gel za jedkanje s fosforno kislino (pribilno 35 %), npr. Scotchbond Universal Echant, ki ga nanesete na sklenko in dentin, in posušite z vodo, potem da reakcija konča.

Temeljito izpirazite z vodo in posušite z zrakom brez primesi vode ali olja ali z vanimi kroglicami; ne susište preveč.

Uporabite običajen gel za jedkanje s fosforno kislino (pribilno 35 %), npr. Scotchbond Universal Echant, ki ga nanesete na pripravljeno in nepravljeno keramiko v sklopu sklenke in dentina, in posušite z vodo, potem da reakcija konča.

Temeljito izpirazite z vodo in posušite z zrakom brez primesi vode ali olja ali z vanimi kroglicami; ne susište preveč.

Uporabite običajen gel za jedkanje s fosforno kislino (pribilno 35 %), npr. Scotchbond Universal Echant, ki ga nanesete na pripravljeno in nepravljeno keramiko v sklopu sklenke in dentina, in posušite z vodo, potem da reakcija konča.

Temeljito izpirazite z vodo in posušite z zrakom brez primesi vode ali olja ali z vanimi kroglicami; ne susište preveč.

Uporabite običajen gel za jedkanje s fosforno kislino (pribilno 35 %), npr. Scotchbond Universal Echant, ki ga nanesete na pripravljeno in nepravljeno keramiko v sklopu sklenke in dentina, in posušite z vodo, potem da reakcija konča.

Temeljito izpirazite z vodo in posušite z zrakom brez primesi vode ali olja ali z vanimi kroglicami; ne susište preveč.

Uporabite običajen gel za jedkanje s fosforno kislino (pribilno 35 %), npr. Scotchbond Universal Echant, ki ga nanesete na pripravljeno in nepravljeno keramiko v sklopu sklenke in dentina, in posušite z vodo, potem da reakcija konča.

Temeljito izpirazite z vodo in posušite z zrakom brez primesi vode ali olja ali z vanimi kroglicami; ne susište preveč.

Uporabite običajen gel za jedkanje s fosforno kislino (pribilno 35 %), npr. Scotchbond Universal Echant, ki ga nanesete na pripravljeno in nepravljeno keramiko v sklopu sklenke in dentina, in posušite z vodo, potem da reakcija konča.

Temeljito izpirazite z vodo in posušite z zrakom brez primesi vode ali olja ali z vanimi kroglicami; ne susište preveč.

Uporabite običajen gel za jedkanje s fosforno kislino (pribilno 35 %), npr. Scotchbond Universal Echant, ki ga nanesete na pripravljeno in nepravljeno keramiko v sklopu sklenke in dentina, in posušite z vodo, potem da reakcija konča.

Temeljito izpirazite z vodo in posušite z zrakom brez primesi vode ali olja ali z vanimi kroglicami; ne susište preveč.

Uporabite običajen gel za jedkanje s fosforno kislino (pribilno 35 %), npr. Scotchbond Universal Echant, ki ga nanesete na pripravljeno in nepravljeno keramiko v sklopu sklenke in dentina, in posušite z vodo, potem da reakcija konča.

Temeljito izpirazite z vodo in posušite z zrakom brez primesi vode ali olja ali z vanimi kroglicami; ne susište preveč.

Uporabite običajen gel za jedkanje s fosforno kislino (pribilno 35 %), npr. Scotchbond Universal Echant, ki ga nanesete na pripravljeno in nepravljeno keramiko v sklopu sklenke in dentina, in posušite z vodo, potem da reakcija konča.

Temeljito izpirazite z vodo in posušite z zrakom brez primesi vode ali olja ali z vanimi kroglicami; ne susište preveč.

Uporabite običajen gel za jedkanje s fosforno kislino (pribilno 35 %), npr. Scotchbond Universal Echant, ki ga nanesete na pripravljeno in nepravljeno keramiko v sklopu sklenke in dentina, in posušite z vodo, potem da reakcija konča.

Temeljito izpirazite z vodo in posušite z zrakom brez primesi vode ali olja ali z vanimi kroglicami; ne susište preveč.

Uporabite običajen gel za jedkanje s fosforno kislino (pribilno 35 %), npr. Scotchbond Universal Echant, ki ga nanesete na pripravljeno in nepravljeno keramiko v sklopu sklenke in dentina, in posušite z vodo, potem da reakcija konča.

Temeljito izpirazite z vodo in posušite z zrakom brez primesi vode ali olja ali z vanimi kroglicami; ne susište preveč.

Uporabite običajen gel za jedkanje s fosforno kislino (pribilno 35 %), npr. Scotchbond Universal Echant, ki ga nanesete na pripravljeno in nepravljeno keramiko v sklopu sklenke in dentina, in posušite z vodo, potem da reakcija konča.

Temeljito izpirazite z vodo in posušite z zrakom brez primesi vode ali olja ali z vanimi kroglicami; ne susište preveč.

Uporabite običajen gel za jedkanje s fosforno kislino (pribilno 35 %), npr. Scotchbond Universal Echant, ki ga nanesete na pripravljeno in nepravljeno keramiko v sklopu sklenke in dentina, in posušite z vodo, potem da reakcija konča.

Temeljito izpirazite z vodo in posušite z zrakom brez primesi vode ali olja ali z vanimi kroglicami; ne susište preveč.

Uporabite običajen gel za jedkanje s fosforno kislino (pribilno 35 %), npr. Scotchbond Universal Echant, ki ga nanesete na pripravljeno in nepravljeno keramiko v sklopu sklenke in dentina, in posušite z vodo, potem da reakcija konča.

Temeljito izpirazite z vodo in posušite z zrakom brez primesi vode ali olja ali z vanimi kroglicami; ne susište preveč.

Uporabite običajen gel za jedkanje s fosforno kislino (pribilno 35 %), npr. Scotchbond Universal Echant, ki ga nanesete na pripravljeno in nepravljeno keramiko v sklopu sklenke in dentina, in posušite z vodo, potem da reakcija konča.

Temeljito izpirazite z vodo in posušite z zrakom brez primesi vode ali olja ali z vanimi kroglicami; ne susište preveč.

Uporabite običajen gel za jedkanje s fosforno kislino (pribilno 35 %), npr. Scotchbond Universal Echant, ki ga nanesete na pripravljeno in nepravljeno keramiko v sklopu sklenke in dentina, in posušite z vodo, potem da reakcija konča.

Temeljito izpirazite z vodo in posušite z zrakom brez primesi vode ali olja ali z vanimi kroglicami; ne susište preveč.

Uporabite običajen gel za jedkanje s fosforno kislino (pribilno 35 %), npr. Scotchbond Universal Echant, ki ga nanesete na pripravljeno in nepravljeno keramiko v sklopu sklenke in dentina, in posušite z vodo, potem da reakcija konča.

Temeljito izpirazite z vodo in posušite z zrakom brez primesi vode ali olja ali z vanimi kroglicami; ne susište preveč.

Uporabite običajen gel za jedkanje s fosforno kislino (pribilno 35 %), npr. Scotchbond Universal Echant, ki ga nanesete na pripravljeno in nepravljeno keramiko v sklopu sklenke in dentina, in posušite z vodo, potem da reakcija konča.

Temeljito izpirazite z vodo in posušite z zrakom brez primesi vode ali olja ali z vanimi kroglicami; ne susište preveč.

Uporabite običajen gel za jedkanje s fosforno kislino (pribilno 35 %), npr. Scotchbond Universal Echant, ki ga nanesete na pripravljeno in nepravljeno keramiko v sklopu sklenke in dentina, in posušite z vodo, potem da reakcija konča.

Temeljito izpirazite z vodo in posušite z zrakom brez primesi vode ali olja ali z vanimi kroglicami; ne susište preveč.

Uporabite običajen gel za jedkanje s fosforno kislino (pribilno 35 %), npr. Scotchbond Universal Echant, ki ga nanesete na pripravljeno in nepravljeno keramiko v sklopu sklenke in dentina, in posušite z vodo, potem da reakcija konča.

Temeljito izpirazite z vodo in posušite z zrakom brez primesi vode ali olja ali z vanimi kroglicami; ne susište preveč.

Uporabite običajen gel za jedkanje s fosforno kislino (pribilno 35 %), npr. Scotchbond Universal Echant, ki ga nanesete na pripravljeno in nepravljeno keramiko v sklopu sklenke in dentina, in posušite z vodo, potem da reakcija konča.

Temeljito izpirazite z vodo in posušite z zrakom brez primesi vode ali olja ali z vanimi kroglicami; ne susište preveč.

Uporabite običajen gel za jedkanje s fosforno kislino (pribilno 35 %), npr. Scotchbond Universal Echant, ki ga nanesete na pripravljeno in nepravljeno keramiko v sklopu sklenke in dentina, in posušite z vodo, potem da reakcija konča.

Temeljito izpirazite z vodo in posušite z zrakom brez primesi vode ali olja ali z vanimi kroglicami; ne susište preveč.

Uporabite običajen gel za jedkanje s fosforno kislino (pribilno 35 %), npr. Scotchbond Universal Echant, ki ga nanesete na pripravljeno in nepravljeno keramiko v sklopu sklenke in dentina, in posušite z vodo, potem da reakcija konča.

Temeljito izpirazite z vodo in posušite z zrakom brez primesi vode ali olja ali z vanimi kroglicami; ne susište preveč.

Uporabite običajen gel za jedkanje s fosforno kislino (pribilno 35 %), npr. Scotchbond Universal Echant, ki ga nanesete na pripravljeno in nepravljeno keramiko v sklopu sklenke in dentina, in posušite z vodo, potem da reakcija konča.

Temeljito izpirazite z vodo in posušite z zrakom brez primesi vode ali olja ali z vanimi kroglicami; ne susište preveč.

Uporabite običajen gel za jedkanje s fosforno kislino (pribilno 35 %), npr. Scotchbond Universal Echant, ki ga nanesete na pripravljeno in nepravljeno keramiko v sklopu sklenke in dentina, in posušite z vodo, potem da reakcija konča.

Temeljito izpirazite z vodo in posušite z zrakom brez primesi vode ali olja ali z vanimi kroglicami; ne susište preveč.

Uporabite običajen gel za jedkanje s fosforno kislino (pribilno 35 %), npr. Scotchbond Universal Echant, ki ga nanesete na pripravljeno in nepravljeno keramiko v sklopu sklenke in dentina, in posušite z vodo, potem da reakcija konča.

Temeljito izpirazite z vodo in posušite z zrakom brez primesi vode ali olja ali z vanimi kroglicami; ne susište preveč.

Uporabite običajen gel za jedkanje s fosforno kislino (pribilno 35 %), npr. Scotchbond Universal Echant, ki ga nanesete na pripravljeno in nepravljeno keramiko v sklopu sklenke in dentina, in posušite z vodo, potem da reakcija konča.

Temeljito izpirazite z vodo in posušite z zrakom brez primesi vode ali olja ali z vanimi kroglicami; ne susište preveč.

Uporabite običajen gel za jedkanje s fosforno kislino (pribilno 35 %), npr. Scotchbond Universal Echant, ki ga nanesete na pripravljeno in nepravljeno keramiko v sklopu sklenke in dentina, in posušite z vodo, potem da reakcija konča.

Temeljito izpirazite z vodo in posušite z zrakom brez primesi vode ali olja ali z vanimi kroglicami; ne susište preveč.

Uporabite običajen gel za jedkanje s fosforno kislino (pribilno 35 %), npr. Scotchbond Universal Echant, ki ga nanesete na pripravljeno in nepravljeno keramiko v sklopu sklenke in dentina, in posušite z vodo, potem da reakcija konča.

