

Surgical Site Infection Risk Reduction Solutions

When it comes to helping ensure patient safety, we're all on the same team.

Protecting patients along the surgical journey is a clinical imperative that depends on the seamless interaction and dedication of professionals like you every step of the way.

Surgical site infections take a tremendous toll on patients, health care providers and facilities. 3M can help you mitigate risk of SSIs and transform patient outcomes for the better.



Surgical site infections: Seeing the big picture.

While progress has been made to reduce SSIs, this is a complication that still occurs too often and may be preventable in up to 60% of cases.¹

SSIs are associated with physical and emotional trauma for patients, increased lengths of stay, resource consumption, and readmission rates. They also represent significant financial burden to health systems as they may not be reimbursable by payors and may increase litigation risk.²



SSI risk factor equation

Reducing a patient's microbial load and improving the resistance of the host are critical to reducing the risk of SSIs.

Every touchpoint matters.

A key to reducing the risk of SSIs is following evidenced based best practice guidelines as recommended by AAMI, AORN, CDC, ERAS, NICE and ACS NSQIP.



SSIs for reporting and practice assessment.^{9,10}
Patients and care partners should also be educated on the risk of SSIs.

Surgical Preparation



Patient Preparation Solution

Take control with a comprehensive approach.

An aging population, a rise in comorbidities, and the emergence of antibiotic resistant bacteria are key factors contributing to the risk of infection in the patient population undergoing surgery.

The overall incidence of infection in surgery is low, but when infections occur, they can be devastating and costly. It is important to take every step possible to help reduce the risk factors of infection for your patients.

Understanding the factors that impact SSIs.

Patient preparation can help reduce the amount of bacteria from the patient's own skin with the right antiseptic products and approach.



Temperature Management

Pre-warm, monitor, maintain.

The difference between a positive patient outcome and a complicated recovery can be a matter of degrees. Unintended perioperative hypothermia is a frequent, yet preventable, complication of surgery. It can increase the rate of wound infection (SSI)^{11,12}, extend recovery time¹³ and length of stay¹⁴, and increase mortality rates¹⁴.

However, hypothermia can be prevented when temperature is monitored and an active warming measure is instituted throughout the perioperative process — starting before the induction of anesthesia.

The 3M[™] Bair Hugger[™] normothermia system provides a comprehensive solution that works seamlessly throughout the perioperative process to effectively and efficiently measure and manage patient temperature.



From noninvasive temperature monitoring and comforting warming gowns to the most extensive array of unique warming blankets, the 3M[™] Bair Hugger[™] normothermia system offers a complete temperature management portfolio.





Surgical Incision Management

Get patients back to daily living.

Surgery has a unique set of challenges; today's environment of value-based care requires providers to constantly refine their practice while continually facing new hurdles.

When surgical site complications occur, like those listed below, they can lead to costly medical interventions (emergency department visits, readmissions, and revisions) and subpar patient experiences.

- Surgical Site Infections
- Necrosis
- Surgical Wound Dehiscence
- EdemaPain

- SeromasHematomas

Meeting the benchmarks of value-based care while catering to patient needs is more difficult than ever. Healthcare organizations need new tools to support surgeons and improve the patient's healing journey.





Sterilization and Cleaning Monitoring

Simplify. Standardize. Streamline.

A unified force to help reduce patient risks. Patients exposed to contaminated instruments are at risk to develop surgical site infections. This exposure risk, plus the rise of antibiotic-resistant bacteria and hardto-clean medical instruments and devices—makes the importance of sterilization and cleaning monitoring especially critical.

C 4hours of lost time

During a recall, tracking down impacted instruments can take as long as 4 hours.¹⁵



Sterilization assurance For all three major modalities: Steam, VH2O2, and EtO



Cleaning monitoring ATP monitoring for endoscopes and high-touch patient surfaces



Digital monitoring and tracking Real-time data accessibility and electronic record-keeping

3M is your team behind the scenes, partnering with clinicians to improve patient outcomes.

3M is a global leader in manufacturing, technology, and lean six sigma processes, and we appreciate the commitment from your facility and staff to reducing process variation and improving patient outcomes. The 3M[™] SSI Risk Reduction solution portfolio allows clinicians to advance quality of care, optimize product utilization, and streamline workflow. 3M can work with your team on implementing recommendations based on industry guidelines using a variety of tools:

- Competency checklists and protocols
- On-site education program
- Implementation of guideline-compliant protocols
- On-line continuing education through the 3M[™] Health Care Academy

Discover how, together you and 3M can help raise the standard of patient care. For more information, contact your 3M representative.

¹Anderson, D., Podgorny, K., Berríos-Torres, S., Bratzler, D., Dellinger, E., Greene, L., . . . Kaye, K. (2014). Strategies to Prevent Surgical Site Infections in Acute Care Hospitals: 2014 Update. *Infection Control & Hospital Epidemiology*, *35*(6), 605-627. doi:10.1086/676022. ²US Department of Health and Human Services. Hospital-Acquired Conditions and Present on Admission Indicator Reporting Provision. ICN 901046.https://www.cms.gov/Outreach-and-Education/Medicare-Learning-Network-MLN/MLNProducts/downloads/ wPOAFactSheet

³Mangram AJ, Horan TC, Pearson ML, Silver LC, Jarvis WR. Guideline for prevention of surgical site infection, 1999. Infect Control Hosp Epidemiol. 1999; 20:24778.

⁴NICE (2019) NICE guideline 125. Surgical site infections: prevention and treatment nice.org.uk/guidance/ng125Centers for Disease Control and Prevention, "Guideline for Prevention of Surgical Site Infections," JAMA Surg. doi:10.1001/jamasurg.2017.0904. ⁵Centers for Disease Control and Prevention, "Guideline for Prevention of Surgical Site Infections," JAMA Surg. doi: 10.1001/jamasurg.2017.0904.

⁶Strategies to Prevent Hospital-onset Staphylococcus aureus Bloodstream Infections in Acute Care Facilities Website. https://www.cdc.gov/hai/prevent/staph-prevention-strategies Reviewed March 1,2019. Accessed August 19,2019.

⁷Kurz A, Sessler DI, Lenhardt R. Perioperative normothermia to reduce the incidence of surgical-wound infection and shorten hospitalization. Study of Wound Infection and Temperature Group. N Engl J Med. 1996;334(19):1209-1215.

⁸AORN. Guideline for Preoperative Environment of Care, Environmental Cleaning and Transmission-based precautions. Guidelines for Perioperative Practices. Denver, Colorado: AORN, Inc. 2020.

⁹DH (2010) The Health and Social Care Act 2008.Code of Practice on the prevention and control of infections and related guidance. ¹⁰Public Health England (June 2013). Protocol for the Surveillance of Surgical Site Infection, Surgical Site Infection Surveillance Service Version 6.

¹¹Kurz A, Sessler DI, et al. Perioperative Normothermia to Reduce the Incidence of Surgical-Wound Infection and Shorten Hospitalization. New Engl J Med. 1996;334:1209-1215.

¹²Melling AC, Ali B, Scott EM, Leaper DJ. Effects of preoperative warming on the incidence of wound infection after a clean surgery: a randomized controlled trial. Lancet. 2001;358(9285):876-880.

¹³Lenhardt R, Marker E, Goll V, et al. Mild intraoperative hypothermia prolongs postanesthetic recovery

Anesthesiology.1997;87(6):1318-1323.

¹⁴Bush H Jr., Hydo J, Fischer E, et al. Hypothermia during elective abdominal aortic aneurysm repair: The high price of avoidable morbidity. J Vasc Surg. 1995;21(3): 392-402.

¹⁵Wong, Janice, and Kathleen Joy Khu. Delays in the Operating Room: Signs of an Imperfect System. Canadian Journal of Surgery June 53.3 (2010): 189-95.

