

Help protect your patients with 3M™ Prevena™ Therapy.

Implement Proactive Risk Management (PRM) with Prevena Therapy.

Prevena Therapy can benefit surgical patients — choosing Prevena Therapy can help you manage and protect the incision in patients at risk of postoperative complications. By implementing PRM, you can use procedural and patient risk stratification to help protect your high-risk patients.

Surgical Site Complications (SSCs) are not only costly, but they can lead to negative impacts on patient recovery.

Surgical Site Infections (SSIs) occur in 2%–5% of all inpatients.¹

Patients who develop an SSI are approximately 5x more likely to be readmitted.²

A single SSI can increase a patient hospital stay by up to 11 days.³

Prevena Therapy has been shown to help reduce the risk of SSCs and overall cost of care.^{4,5}

Prevena Therapy has demonstrated outcomes across multiple specialties, including plastic, vascular, cardiothoracic, spine, orthopedic and general surgery.⁶ Data from a multicentre randomized controlled trial and health economic analysis showed that 3M™ Prevena™ Therapy significantly reduced the risk of 90-day surgical site complications (SSCs),⁴ readmissions,⁴ and surgical site management costs⁵ vs. silver-impregnated dressings.



4x
reduction
in SSCs.¹⁴

3.4% (5/147) Prevena Therapy vs. 14.3% (21/147) SOC (p=0.0013)[†]



3x
reduction
in readmission rates.¹⁴

3.4% (5/147) Prevena Therapy vs. 10.2% (15/147) SOC (p=0.0208)[†]



1.9x
reduction
in per-patient cost of care.⁵

\$1,047 Prevena Therapy vs. \$2,036 SOC

* The effectiveness of Prevena Therapy in reducing the incidence of SSIs and seroma in all surgical procedures and populations has not been demonstrated. See full indications for use and limitations at mykci.com.

† Calculation(s) are derived based on relative patient group incidence rate reported in this study.

‡ Statistically significant (p<0.05)

For additional information specific to your specialty visit 3M.ca/Prevena

The 3M™ Prevena™ Incision Management System is intended to manage the environment of surgical incisions and surrounding intact skin in patients at risk for developing post-operative complications, such as infection, by maintaining a closed environment via the application of a negative pressure wound therapy system to the incision. The 3M™ Prevena™ Dressing skin interface layer with silver reduces microbial colonization in the fabric.

Available in Canada from your authorized 3M+KCI distributors.
KCI USA Inc., a 3M Company
KCI owned and operated by 3M Company

KCI Medical Canada Inc.
75 Courtneypark Dr W, Unit 4
Mississauga, ON
L5W 0E3

KCI USA Inc.
12930 IH 10 West
San Antonio, TX
78249
USA

1. Anderson, DJ, et al. Strategies to Prevent Surgical Site Infections in Acute Care Hospitals: 2014 Update. *Infect Control Hosp Epidemiol.* 2014;35(6):605-627. doi: 10.1086/676022.

2. Canadian Surgical Site Infection Prevention Audit Month Report. Retrieved from <http://www.patientsafetyinstitute.ca/en/toolsResources/Pages/SSI-Audit-Recap-Report-2016-12.aspx>

3. Canadian Patient Safety Institute. Canadian Surgical Site Infection Prevention Audit Month. February 2016. CPSI; 2016. http://www.patientsafetyinstitute.ca/en/toolsResources/Documents/SSI Audit 2016_Recap Report EN.pdf

4. Higuera-Rueda C, et al. The Effectiveness of Closed Incision Negative Pressure Therapy versus Silver-Impregnated Dressings in Mitigating Surgical Site Complications in High-Risk Patients after Revision Knee Arthroplasty: The PROMISES Randomized Controlled Trial. *J Arthroplasty.* 2021;36(7S):S295-S302.e14. doi: 10.1016/j.arth.2021.02.076.

5. Cooper HJ, Bongards C, Silverman RP. Cost-effectiveness of closed incision negative pressure therapy for surgical site management after revision total knee arthroplasty: Secondary analysis of a randomized clinical trial. Presented at: American Association of Hip and Knee Surgeons Annual Meeting, November 11-14, 2021, Dallas, Texas.

6. Singh DP, Gabriel A, Parvizi J, Gardner MJ, D'Agostino R Jr. Meta-Analysis of Comparative Trials Evaluating a Single-Use Closed-Incision Negative-Pressure Therapy System. *Plast Reconstr Surg.* 2019 Jan;143(1S Management of Surgical Incisions Utilizing Closed-Incision Negative-Pressure Therapy):41S-46S.

NOTE: Specific indications, limitations, contraindications, warnings, precautions and safety information exist for these products and therapies. Please consult a clinician and product instructions for use prior to application. This material is intended for healthcare professionals only.

3M, and Prevena are trademarks of 3M. Used under license in Canada. © 2022, 3M. All rights reserved. 2208-24635-a-E