

Before you select a surgical patient skin preparation, consider which prep makes the most sense for the type of procedure you are doing.

ChloraPrep® Patient Preoperative Skin Preparation

2% Chlorhexidine Gluconate (CHG) & 70% Isopropyl Alcohol (IPA)

3M™ DuraPrep™ Surgical Solution

(lodine Povacrylex [0.7% available iodine] and Isopropyl Alcohol, 74% w/w) Patient Preoperative Skin Preparation


Product Cost


\$ per 26 mL applicator*

*ChloraPrep solution brochure: "Surgical site infections: the economic impact" \$ per 26 mL applicator**

**Data on file.

Coverage Area





Required Application Technique

You must scrub. Per manufacturer's instructions for use: "Dry Site (such as abdomen or arm) Use repeated back-and-forth strokes of the sponge for approximately 30 seconds. Completely wet the treatment area with antiseptic.



Moist Site (such as the inguinal fold) Use repeated back-and-forth strokes of the sponge for approximately 2 minutes. Completely wet the treatment area with antiseptic."

You can paint. Per manufacturer's instructions for use: "Paint a single, uniform application and do not go back over areas already prepped."



Guidelines Met

CDC AORN

CDC AORN

3M Infection Prevention Solutions

Innovation
On A Mission

