

For use as a surgical hand antiseptic

3M

Avagard™

NDC 17518-051-01

(Chlorhexidine Gluconate 1% Solution and Ethyl Alcohol 61% w/w)
*surgical and healthcare personnel hand antiseptic
with moisturizers*

WARNING: FLAMMABLE. DO NOT TOUCH THE EYE WITH HANDS THAT HAVE BEEN TREATED WITH THIS PREPARATION. DO NOT USE • if you are allergic to chlorhexidine gluconate or any other ingredient in this preparation • routinely if you have wounds which involve more than the superficial layers of the skin.

Stop use and ask a doctor if irritation, sensitization or allergic reaction occurs. These may be signs of a serious condition.

Read Drug Facts Panel before use.

Directions: Surgical hand antiseptic

- apply to clean, dry hands and nails. For the first use of each day, clean under nails with a nail stick.
- dispense one pump (2 mL) into the palm of one hand
- dip the fingertips of the opposite hand into the lotion and work it under the nails
- spread the remaining lotion over the hand and up to just above the elbow
- using another 2 mL of lotion, repeat with the other hand
- dispense another 2 mL of lotion into either hand and reapply to all aspects of both hands up to the wrist
- allow to dry before donning gloves

16 fl oz • 500 mL

Peel off to view directions for Use as a Healthcare Personnel Hand Antiseptic

U.S. Patent Nos. 5,799,841; 5,897,031; 6,090,395; 6,534,069; 6,623,744; 7,081,246; 7,566,460

Drug Facts

Active ingredients

Chlorhexidine gluconate 1% w/w.....	Antiseptic
Ethyl alcohol 61% w/w.....	Antiseptic

Purpose

Uses • **surgical hand antiseptic:** significantly reduces the number of micro-organisms on the hands and forearms prior to surgery or patient care

- **healthcare personnel hand antiseptic:** handwash to help reduce bacteria that potentially can cause disease • recommended for repeated use

Warnings

Flammable, keep away from fire or flame. For external use only

Do not use if you are allergic to chlorhexidine gluconate or any other ingredients in this preparation

When using this product • do not touch the eye with hands that have been treated with this preparation • keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye. • if contact occurs, rinse with cold water right away • do not use routinely if you have wounds which involve more than the superficial layers of the skin

Stop use and ask a doctor if irritation, sensitization or allergic reaction occurs. These may be signs of a serious condition

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. ▶

Drug Facts (continued)

Directions Use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.

Surgical hand antiseptic • apply to clean, dry hands and nails. For the first use of each day, clean under nails with a nail stick. • dispense one pump (2 mL) into the palm of one hand • dip the fingertips of the opposite hand into the lotion and work it under the nails • spread the remaining lotion over the hand and up to just above the elbow • using another 2 mL of lotion, repeat with the other hand

- dispense another 2 mL of lotion into either hand; reapply to all aspects of both hands up to the wrists • allow to dry before donning gloves

Healthcare personnel hand antiseptic • apply to clean, dry hands and nails • dispense one pump (2 mL) into the palm of one hand • paying particular attention to the spaces between the fingers and under fingernails, apply the lotion evenly to cover both hands up to the wrists

- allow to dry without wiping

Other information store at 20-25°C (68-77°F)

Inactive ingredients beheneth 10, behenyl alcohol, C20-40 pareth-24, cetyl palmitate, diisopropyl dimer dilinoleate, dimethicone, glycerin, polyethylene glycol, squalane, water

Questions? call 1-800-228-3957

Made in U.S.A. for 3M Health Care
2510 Conway Ave., St. Paul, MN 55144
www.3M.com

The shape and colors of the bottle and wall bracket are trademarks of 3M.
© 2012, 3M. All rights reserved.

REF
9200

MAL10110039XC

For use as a healthcare personnel hand antiseptic



Avagard™

NDC 17518-051-01

(Chlorhexidine Gluconate 1% Solution and Ethyl Alcohol 61% w/w)
*surgical and healthcare personnel hand antiseptic
with moisturizers*

WARNING: FLAMMABLE. DO NOT TOUCH THE EYE WITH HANDS THAT HAVE BEEN TREATED WITH THIS PREPARATION. DO NOT USE • if you are allergic to chlorhexidine gluconate or any other ingredient in this preparation • routinely if you have wounds which involve more than the superficial layers of the skin.

Stop use and ask a doctor if irritation, sensitization or allergic reaction occurs. These may be signs of a serious condition.

Read Drug Facts Panel before use.

Directions: *Healthcare personnel hand antiseptic*

- apply to clean, dry hands and nails
- dispense one pump (2 mL) into the palm of one hand
- paying particular attention to the spaces between the fingers and under fingernails, apply the lotion evenly to cover both hands up to the wrists
- allow to dry without wiping

16 fl oz • 500 mL

U.S. Patent Nos. 5,799,841; 5,897,031; 6,090,395; 6,534,069; 6,623,744; 7,081,246; 7,566,460

Drug Facts

Active ingredients

Chlorhexidine gluconate 1% w/w.....	Antiseptic
Ethyl alcohol 61% w/w.....	Antiseptic

Purpose

Uses • **surgical hand antiseptic:** significantly reduces the number of micro-organisms on the hands and forearms prior to surgery or patient care

- **healthcare personnel hand antiseptic:** handwash to help reduce bacteria that potentially can cause disease • recommended for repeated use

Warnings

Flammable, keep away from fire or flame. For external use only

Do not use if you are allergic to chlorhexidine gluconate or any other ingredients in this preparation

When using this product • do not touch the eye with hands that have been treated with this preparation • keep out of eyes, ears, and mouth. May cause serious and permanent eye injury if permitted to enter and remain in the eye. • if contact occurs, rinse with cold water right away • do not use routinely if you have wounds which involve more than the superficial layers of the skin

Stop use and ask a doctor if irritation, sensitization or allergic reaction occurs. These may be signs of a serious condition

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. ▶

Drug Facts (continued)

Directions Use with care in premature infants or infants under 2 months of age. These products may cause irritation or chemical burns.

Surgical hand antiseptic • apply to clean, dry hands and nails. For the first use of each day, clean under nails with a nail stick. • dispense one pump (2 mL) into the palm of one hand • dip the fingertips of the opposite hand into the lotion and work it under the nails • spread the remaining lotion over the hand and up to just above the elbow • using another 2 mL of lotion, repeat with the other hand

- dispense another 2 mL of lotion into either hand; reapply to all aspects of both hands up to the wrists • allow to dry before donning gloves

Healthcare personnel hand antiseptic • apply to clean, dry hands and nails • dispense one pump (2 mL) into the palm of one hand • paying particular attention to the spaces between the fingers and under fingernails, apply the lotion evenly to cover both hands up to the wrists

- allow to dry without wiping

Other information store at 20-25°C (68-77°F)

Inactive ingredients beheneth 10, behenyl alcohol, C20-40 pareth-24, cetyl palmitate, diisopropyl dimer dilinoleate, dimethicone, glycerin, polyethylene glycol, squalane, water

Questions? call 1-800-228-3957

Made in U.S.A. for 3M Health Care
2510 Conway Ave., St. Paul, MN 55144
www.3M.com

The shape and colors of the bottle and wall bracket are trademarks of 3M.
© 2012, 3M. All rights reserved.

REF
9200

MAL10110039XC